

# Wound Care Cream

Now available from Letco Medical!

new

## The optimal healing environment for wound-damaged skin.

Letco Medical Wound Care base is a polyethylene glycol (PEG) gel clinically shown to promote the optimal healing environment for immediate and ongoing wound management. Our advanced base quiets the wound area and cultivates clean, moist conditions to encourage faster and more noticeable healing.

Polyethylene glycol-based preparations are preferred protocol for aiding in the wound healing process, and in gel form provide distinct advantages. The other key ingredient in our base, organic meadowsweet extract, or meadowort, is noted for its potential to inhibit germs. It has also exhibited excellent anti-inflammatory and healing properties. Our base can further enhance the healing process by delivering water compatible APIs to sensitive wound sites.

Letco Medical's commitment to quality is clearly evident in this water-soluble product. It is prepared with your patients in mind, designed to provide them with the ideal combination of ingredients for superior wound care.



QUALITY COMPOUNDING STARTS HERE

# Wound Care Cream

## Polyethylene glycol (PEG) gel creates the perfect healing environment

- Hydrophilic base won't evaporate and maintains consistent moisture content
- Literature suggests PEG-based occlusive dressings may aid in the wound healing process<sup>1-4</sup>
- PEG gel migrates to deeper wound areas
- Gel form is more consumer friendly

## Organic meadowsweet extract

- Also known as meadow, its high content of phenolic compounds has shown the potential to provide germicidal inhibitions<sup>5-6</sup>
- Salicylates and flavonoids found in the extract exhibited anti-inflammatory and healing properties<sup>7-10</sup>

## The ideal benefits for successful wound treatment

- Maintains a moist wound site and the optimal healing environment
- Occlusive and adherent
- Water-washable for easy cleaning and debridement
- Gel base is non-greasy, quick drying, colorless, odorless, and water-soluble
- Invisible when dry
- Safe for sensitive skin
- pH: 5.8
- Specific Gravity: 0.96 g/cm<sup>3</sup>

## Formulated to excel on wounds and more

- Burns
- Cuts and abrasions
- Dermatological applications
- Sores
- Ulcers

**Note:** For external use only. Medical research references available upon request. Use under medical supervision. Letco Wound Care Cream should not be used in conditions where absorption of large quantities of polyethylene glycol is possible, for example on extensive burn areas and large surface areas, especially if there is evidence of moderate or severe renal impairment.

Item No.	Size
693800	100g
693801	500g
693802	2.5kg

## Sample Formulas:

Gentamicin 1 mg/g	Itraconazole 5%
Clindamycin 1mg/g	Fluticasone Propionate 1%
Polymyxin 2,000 u/g	Letco Wound Care Cream qs 100%
Letco Wound Care Cream qs 100%	.....
Morphine Sulfate 1%	Lidocaine 5%
Letco Wound Care Cream qs 100%	Urea 30%
	Letco Wound Care Cream qs 100%



## References

1. Mogosanu GD, Grumezescu AM. Natural and synthetic polymers for wounds and burns dressing. *Int J Pharm.* 2014 Mar; 463(2): 127-36.
2. Mhessn R.J. et al. Preparation of Tannin Based Hydrogel for Biological Application. *Iranian Journal of Pharmaceutical Sciences Autumn* 2010; 6(4): 263-268.
3. Anumolu SS. et al. Doxycycline loaded poly(ethylene glycol) hydrogels for healing vesicant-induced ocular wounds. *Biomaterials.* 2010 Feb;31(5):964-74.
4. Cho CY, Lo JS. Excision and repair: dressing the part. *Dermatol Clin* 1998; 16:25-47.
5. Churin AA et al. Effect of Filipendula ulmaria extract on immune system of CBA/ CaLac and C57B/6 Mice. *Eksp Klin Farmakol.* 2008 Sep-Oct;71(5):32-6.
6. Rauha JP et al. Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. *Int J Food Microbiol.* 2000 May 25;56(1):3-12.
7. Liapina LA, Koval'chuk GA. A comparative study of the action on the hemostatic system of extracts from the flowers and seeds of the meadowsweet (*Filipendula ulmaria* (L.) Maxim. *Izv Akad Nauk Ser Biol.* 1993 Jul-Aug;(4):625-8.
8. Nitta Y, et al. Inhibitory activity of Filipendula ulmaria constituents on recombinant human histidine decarboxylase. *Food Chem.* 2013 Jun 1;138(2-3):1551-6.
9. Drummond EM, et al. Inhibition of proinflammatory biomarkers in THP1 macrophages by polyphenols derived from chamomile, meadowsweet and willow bark. *Phytother Res.* 2013 Apr; 27(4):588-94
10. Nesterova IuV, et al. Evaluation of anti-inflammatory activity of extracts from Siberian plants. *Vestn Ross Akad Med Nauk.* 2009;(11):30-4.



QUALITY COMPOUNDING STARTS HERE

800.239.5288 • www.letcomedical.com